

2017年 英文誌掲載論文

Twelve-year trends of increasing overweight and obesity in patients with diabetes: the Shiga Diabetes Clinical Survey.	Miyazawa I, Kadota A, Miura K, Okamoto M, Nakamura T, Ikai T, Maegawa H, Ohnishi A.	Endocr J.
Safety and efficacy of ipragliflozin in elderly versus non-elderly Japanese patients with type 2 diabetes mellitus: a subgroup analysis of the STELLA-LONG TERM study.	Maegawa H, Tobe K, Tabuchi H, Nakamura I, Uno S.	Expert Opin Pharmacother.
Effects of Pemafibrate, a Novel Selective PPAR α Modulator, on Lipid and Glucose Metabolism in Patients With Type 2 Diabetes and Hypertriglyceridemia: A Randomized, Double-Blind, Placebo-Controlled, Phase 3 Trial.	Araki E, Yamashita S, Arai H, Yokote K, Satoh J, Inoguchi T, Nakamura J, Maegawa H, Yoshioka N, Tanizawa Y, Watada H, Suganami H, Ishibashi S.	Diabetes Care.
Lack of O-GlcNAcylation enhances exercise-dependent glucose utilization potentially through AMP-activated protein kinase activation in skeletal muscle.	Murata K, Morino K, Ida S, Ohashi N, Lemecha M, Park SY, Ishikado A, Kume S, Choi CS, Sekine O, Ugi S, Maegawa H.	Biochem Biophys Res Commun.
Definitive diagnosis of mandibular hypoplasia, deafness, progeroid features and lipodystrophy (MDPL) syndrome caused by a recurrent de novo mutation in the POLD1 gene.	Sasaki H, Yanagi K, Ugi S, Kobayashi K, Ohkubo K, Tajiri Y, Maegawa H, Kashiwagi A, Kaname T.	Endocr J
Safety and efficacy of ipragliflozin in Japanese patients with type 2 diabetes in real-world clinical practice: interim results of the STELLA-LONG TERM post-marketing surveillance study.	Nakamura I, Maegawa H, Tobe K, Tabuchi H, Uno S.	Expert Opin Pharmacother.
Overexpression of acetyl CoA carboxylase β exacerbates podocyte injury in the kidney of streptozotocin-induced diabetic mice.	Tanaka Y, Kume S, Maeda S, Osawa N, Takeda N, Chin-Kanasaki M, Isshiki K, Ugi S, Oshima I, Uzu T, Maegawa H, Araki SI.	Biochem Biophys Res Commun.
Impact of obesity on annual medical expenditures and diabetes care in Japanese patients with type 2 diabetes mellitus.	Kusunoki-Tsuji C, Araki SI, Kume S, Chin-Kanasaki M, Osawa N, Morino K, Sekine O, Ugi S, Kashiwagi A, Maegawa H.	J Diabetes Investig.
Effect of an intensified multifactorial intervention on cardiovascular outcomes and mortality in type 2 diabetes (J-DOIT3): an open-label, randomised controlled trial.	Ueki K, Sasako T, Okazaki Y, Kato M, Okahata S, Katsuyama H, Haraguchi M, Morita A, Ohashi K, Hara K, Morise A, Izumi K, Ishizuka N, Ohashi Y, Noda M, Kadowaki T; J-DOIT3 Study Group.	Lancet Diabetes Endocrinol.
Pivotal Role of O-GlcNAc Modification in Cold-Induced Thermogenesis by Brown Adipose Tissue Through Mitochondrial Biogenesis.	Ohashi N, Morino K, Ida S, Sekine O, Lemecha M, Kume S, Park SY, Choi CS, Ugi S, Maegawa H.	Diabetes
Current status of achieving blood pressure target and its clinical correlates in Japanese type 2 diabetes.	Yokoyama H, Araki SI, Kawai K, Hirao K, Kurihara Y, Seino H, Takamura H, Sugimoto H, Okada A, Maegawa H; JDDM study group.	J Diabetes Investig.
N-3 Polyunsaturated Fatty Acids Decrease the Protein Expression of Soluble Epoxide Hydrolase via Oxidative Stress-Induced P38 Kinase in Rat Endothelial Cells.	Okada T, Morino K, Nakagawa F, Tawa M, Kondo K, Sekine O, Imamura T, Okamura T, Ugi S, Maegawa H.	Nutrients.
Fiber-rich diet with brown rice improves endothelial function in type 2 diabetes mellitus: A randomized controlled trial.	Kondo K, Morino K, Nishio Y, Ishikado A, Arima H, Nakao K, Nakagawa F, Nikami F, Sekine O, Nemoto KI, Suwa M, Matsumoto M, Miura K, Makino T, Ugi S, Maegawa H.	PLoS One.
Diverse metabolic effects of O-GlcNAcylation in the pancreas but limited effects in insulin-sensitive organs in mice.	Ida S, Morino K, Sekine O, Ohashi N, Kume S, Chano T, Iwasaki K, Harada N, Inagaki N, Ugi S, Maegawa H.	Diabetologia.
Acute Effect of Metformin on Postprandial Hypertriglyceridemia through Delayed Gastric Emptying.	Sato D, Morino K, Nakagawa F, Murata K, Sekine O, Beppu F, Gotoh N, Ugi S, Maegawa H.	Int J Mol Sci.