

## 2021年 英文誌掲載論文

Combination of disease duration-to-age at diagnosis and hemoglobin A1c-to-serum C-peptide reactivity ratios predicts patient response to glucose-lowering medication in type 2 diabetes: A retrospective cohort study across Japan (JDDM59)	Kanatsuka A, Sato Y, Higashi Y, Goto Y, Kawai K, Maegawa H; Japan Diabetes Data Management Study Group (JDDM).	J Diabetes Investig.
Relationship between Kidney Function and Subclinical Atherosclerosis Progression Evaluated by Coronary Artery Calcification	Namuun Ganbaatar, Aya Kadota, Takashi Hisamatsu, Shinichi Araki, Shinji Kume, Akira Fujiyoshi, Sayaka Kadowaki, Sayuki Torii, Keiko Kondo, Hiroyoshi Segawa, Ebtehal Salman, Itsuko Miyazawa, Takashi Yamamoto, Yoshihisa Nakagawa, Hiroshi Maegawa, Katsuyuki Miura, Hirotsugu Ueshima, for the SESSA Research Group	J Atheroscler Thromb.
Correction to: Safety and Effectiveness of Ipragliflozin in Elderly Versus Non-elderly Japanese Patients with Type 2 Diabetes: Subgroup Analysis of STELLA-LONG TERM	Nakamura I, Maegawa H, Tobe K, Uno S.	Diabetes Ther.
Comparing Associations of Dietary Energy Density and Energy Intake, Macronutrients with Obesity in Patients with Type 2 Diabetes (JDDM 63)	Takeda Y, Fujihara K, Nedachi R, Ikeda I, Morikawa SY, Hatta M, Horikawa C, Kato M, Kato N, Yokoyama H, Kurihara Y, Miyazawa K, Maegawa H, Sone H.	Nutrients.
Glycemic control and number of natural teeth: analysis of cross-sectional Japanese employment-based dental insurance claims and medical check-up data	Harada K, Morino K, Ishikawa M, Miyazawa I, Yasuda T, Hayashi M, Ishikado A, Maegawa H.	Diabetol Int.
Effect of ipragliflozin on liver function in Japanese type 2 diabetes mellitus patients: subgroup analysis of a 3-year post-marketing surveillance study (STELLA-LONG TERM)	Tobe K, Maegawa H, Nakamura I, Uno S.	Endocr J.
Correction to: Safety and Effectiveness of Ipragliflozin for Type 2 Diabetes in Japan: 12-Month Interim Results of the STELLA-LONG TERM Post-Marketing Surveillance Study	Nakamura I, Maegawa H, Tobe K, Uno S.	Adv Ther.
Patient characteristics associated with improvement in glycemic control following addition of an oral antidiabetic drug to DPP-4 inhibitor monotherapy in Japanese patients with type 2 diabetes mellitus (JDDM 60)	Kawai K, Eiki JI, Kanatsuka A, Motohashi S, Wakana A, Hayashi A, Iglay K, Yamazaki K, Tokita S, Maegawa H; Japan Diabetes Clinical Data Management Study Group (JDDM).	Diabetol Int.
Correction to: Safety and Effectiveness of Ipragliflozin in Elderly Versus Non-elderly Japanese Patients with Type 2 Diabetes: Subgroup Analysis of STELLA-LONG TERM	Nakamura I, Maegawa H, Tobe K, Uno S.	Diabetes Ther.
Malfunctioning CD106-positive, short-term hematopoietic stem cells trigger diabetic neuropathy in mice by cell fusion	Katagi M, Terashima T, Ohashi N, Nakae Y, Yamada A, Nakagawa T, Miyazawa I, Maegawa H, Okano J, Suzuki Y, Fujino K, Eguchi Y, Kojima H.	Commun Biol.
Cardio- and reno-protective effects of dipeptidyl peptidase III in diabetic mice	Komeno M, Pang X, Shimizu A, Molla MR, Yasuda-Yamahara M, Kume S, Rahman NIA, Soh JEC, Nguyen LKC, Ahmat Amin MKB, Kokami N, Sato A, Asano Y, Maegawa H, Ogita H.	J Biol Chem.
Real-world evidence for long-term safety and effectiveness of ipragliflozin in treatment-naïve versus non-naïve Japanese patients with type 2 diabetes mellitus: subgroup analysis of a 3-year post-marketing surveillance study (STELLA-LONG TERM)	Maegawa H, Tobe K, Nakamura I, Uno S.	Diabetol Int.
Clinical inertia in patients with type 2 diabetes treated with oral antidiabetic drugs: Results from a Japanese cohort study (JDDM53)	Maegawa H, Ishigaki Y, Langer J, Saotome-Nakamura A, Andersen M; Japan Diabetes Clinical Data Management (JDDM) Study Group.	J Diabetes Investig.
Targeted deletion of Atg5 in intestinal epithelial cells promotes dextran sodium sulfate-induced colitis	Nishino K, Nishida A, Inatomi O, Imai T, Kume S, Kawahara M, Maegawa H, Andoh A.	J Clin Biochem Nutr
Role of O-linked N-acetylglucosamine in the homeostasis of metabolic organs, and its potential links with diabetes and its complications	Morino K, Maegawa H.	J Diabetes Investig.
Real-World Evidence for Long-Term Safety and Effectiveness of Ipragliflozin in Japanese Patients with Type 2 Diabetes Mellitus: final Results of a 3-Year Post-Marketing Surveillance Study (STELLA-LONG TERM)	Nakamura I, Maegawa H, Tobe K, Uno S.	Expert Opin Pharmacother.
Genome-Wide Association Studies Identify Two Novel Loci Conferring Susceptibility to Diabetic Retinopathy in Japanese Patients with Type 2 Diabetes	Imamura M, Takahashi A, Matsunami M, Horikoshi M, Iwata M, Araki SI, Toyoda M, Susarla G, Ahn J, Park KH, Kong J, Moon S, Sobrin L, Yamauchi T, Tobe K, Maegawa H, Kadowaki T, Maeda S.	Hum Mol Genet.
Roles of mTOR in Diabetic Kidney Disease	Yasuda-Yamahara M, Kume S, Maegawa H.	Antioxidants (Basel).
Machine Learning Approach to Decision Making for Insulin Initiation in Japanese Patients With Type 2 Diabetes (JDDM 58): Model Development and Validation Study	Fujihara K, Matsubayashi Y, Harada Yamada M, Yamamoto M, Iizuka T, Miyamura K, Hasegawa Y, Maegawa H, Kodama S, Yamazaki T, Sone H.	JMIR Med Inform.
MicroRNA-494-3p inhibits formation of fast oxidative muscle fibres by targeting E1A-binding protein p300 in human-induced pluripotent stem cells	Iwasaki H, Ichihara Y, Morino K, Lemecha M, Sugawara L, Sawano T, Miake J, Sakurai H, Nishi E, Maegawa H, Imamura T.	Sci Rep.
Sodium-glucose cotransporter 2 inhibitors represent a paradigm shift in the prevention of heart failure in type 2 diabetes patients	Kashiwagi A, Araki S, Maegawa H.	J Diabetes Investig.
Geometry of Sleeve Gastrectomy Measured by 3D CT Versus Weight Loss: Preliminary Analysis	Yamaguchi T, Yamamoto H, Tomozawa Y, Ugi S, Kaida S, Miyake T, Morino K, Watanabe Y, Maegawa H, Tani M.	World J Surg.
Glucagon-Like Peptide-1 Receptor Agonist Utilization in Type 2 Diabetes in Japan: A Retrospective Database Analysis (JDDM 57)	Ishigaki Y, Strizek A, Aranishi T, Arai N, Imaoka T, Cai Z, Maegawa H.	Diabetes Ther.